

Benghal Dayflower

[*Commelina benghalensis* (L.) Small]

Victor Maddox, Geosystems Research Institute, Mississippi State University

John D. Byrd, Jr., Plant and Soil Sciences, Mississippi State University

Randy Westbrook, Invasive Species Specialist, U.S. Geological Survey, BRD

Row Crop



Fig. 1. The underground stolons of the Benghal dayflower can produce flowers underground.

Fig. 2. Benghal dayflower has simple, opposite leaves. Short hairs grow on the leaves and stems.

Fig. 3. Purple-blue flowers grow in the leaf axils.

Introduction

Problems Caused

Benghal dayflower (*Commelina benghalensis* L.) [Syn. *Commelina benghalensis* L. var. *benghalensis* C.B. Clarke], also known as tropical spiderwort, is a herbaceous perennial first observed in Florida in 1928, then in Georgia in 1967. This Federal noxious weed is native to Africa and tropical Asia and, since its introduction, has become a serious pest in Florida and Georgia. Recently, it was reported in California, Louisiana and North Carolina. In August 2006, it was found in Jackson County, Mississippi. This invasive plant can tolerate a wide range of environmental conditions and can establish dense, monospecific stands, particularly in Roundup Ready® cropping systems.

Regulations

Benghal dayflower is a Federal noxious weed, which means it is a violation of Federal law to transport this plant across a state line. In the southeastern United States, it is listed as a noxious weed in Alabama, Florida, North Carolina, and Mississippi.

Description

Vegetative Growth

Benghal dayflower is an annual or perennial herb with simple, alternate leaves (Figure 2), approximately 2" long and nearly as wide. Leaves and aboveground stems have short hairs (pubescent) and longer red hairs on the leaf sheath and petiole margins. Stems often root at the nodes and purple-blue aerial flowers (chasmogamous) (Figure 3) may be produced in the leaf axils. Underground stolons can produce subterranean flowers (cleistogamous) (Figure 1). Stem cuttings on the soil surface can regenerate easily, although cuttings buried deeper than about 1" fail to regenerate. Broken stems may persist on the soil surface for several weeks or months in low moisture conditions and easily form leaves after moisture becomes available.

Flowering

Both aerial (chasmogamous) (Figure 3) and subterranean (cleistogamous) (Figure 1) flowers can produce seed, although seeds are dimorphic. Aerial seeds are small with five seeds per capsule, while subterranean seeds are large with three seeds per capsule. Both have a rough surface. Plants, flowers, seeds, and chromosome number can be variable. Plants reproduce both sexually and asexually. A single plant can produce 1600 seeds. Fresh, aerial seeds are dormant because of an impermeable seed coat. Dormancy can be broken by scarification. The smaller aerial seeds tend to germinate at shallower soil depths compared to larger subterranean seeds. Seasonal germination can occur over an extended period of time.

Dispersal

Benghal dayflower is most problematic in row crop cropping systems, such as peanuts, soybeans and corn. Dispersal mechanisms may include farm equipment and products. It has been reported in nursery containers, but this seems to be a minor mechanism of dispersal.

Spread By

Benghal dayflower can be spread by farm equipment and products and, to a lesser degree, nursery containers.

Habitat

Benghal dayflower is a terrestrial perennial that occurs in both wet and dry lands, but grows best in moist, highly fertile soils. Because it has exhibited tolerance to Roundup®, it has become particularly troublesome in Roundup Ready® cropping systems in the United States. It is troublesome in cotton, soybeans, peanuts, and, to a lesser extent, corn. However, it can also be a weed in natural areas, roadsides, waste places, along dikes, irrigation ditch banks, field borders, wet pasturelands and gardens. It has also been found in nursery containers, another possible means of movement.

Distribution

U.S. - Benghal dayflower is native to Africa and Tropical Asia, but has become a weed worldwide in the tropics and subtropics. In the United States, it has been reported in Alabama, California, Florida, Georgia, Louisiana, Mississippi, and North Carolina.

Mid-South - Benghal dayflower has been reported in Louisiana, and more recently, Alabama and Mississippi.

Control Methods

Biological - No biological control methods are currently known to be registered or utilized in the United States.

Chemical - Several herbicides, such as MSMA, glyphosate, and 2,4-D provide effective postemergence control when applied to small, actively growing plants. Since Benghal dayflower has shown tolerance to Roundup, caution should be taken regarding application rates and plant maturity. Because of the high germination rates of seed and the length of germination periods, the key to season-long control of Benghal dayflower is to include a herbicide that provides residual preemergence control. Dual Magnum (s-metolachlor) and other chloroacetamide herbicides, such as Lasso (alachlor) and Outlook (dimethanamid) are effective, although Dual Magnum provides the longest residual control. Herbicides frequently used for noncropland weed control are currently being evaluated for control of Benghal dayflower. Check with your local county extension office for additional information.

Mechanical - Hand removal may be possible for very small infestations. Seedlings in moist soil can often be pulled up carefully to ensure complete removal. Early detection and eradication is important, since larger infestations may require broadcast herbicide applications. In addition, control of Benghal dayflower populations can be achieved by moldboard plowing plant material more than 1 inch below the soil surface.

References

Ferrell, J.A., G.E. MacDonald, and B.J. Brecke. 2004. Tropical Spiderwort (*Commelina benghalensis* L.), Identification and Control. SS-AGR-223. University of Florida, Gainesville, FL.

Prostko, E.P., A.S. Culpepper, T.M. Webster, and J.T. Flanders. 2005. Tropical Spiderwort Identification and Control in Georgia Field Crops. Circular 884. University of Georgia, Tifton, GA.

SDA, NRCS. 2007. The PLANTS Database (<http://plants.usda.gov>, 5 September 2007). National Plant Data Center, Baton Rouge, LA 70874-

More Information

Benghal dayflower is in the Commelinaceae Family. There are around 170 species of *Commelina* L. worldwide, most native to Africa. Nine species exist in the United States, with the highest number of species in Florida. *Commelina benghalensis* var. *benghalensis* is widespread worldwide including the United States while *C. benghalensis* var. *hirsuta* is widespread only in Africa. *Commelina benghalensis* var. *benghalensis* is discussed in this factsheet. Other species that occur in the mid-south include Asiatic dayflower (*C. communis* L.), Carolina dayflower (*Commelina caroliniana* Walt.), common dayflower (*C. diffusa* Burm. f. var. *diffusa*), Virginia dayflower (*C. virginica* L.), and white-mouth dayflower (*C. erecta* L.). All have flowers with two to three bright blue petals. However, only Benghal dayflower has subterranean flowers (Figure 1). Both Benghal dayflower and Virginia dayflower may produce underground stems and red hairs, but blue petals and proportionally longer leaves on Virginia dayflower help differentiate it from Benghal dayflower which has broader leaves and purple-blue petals. Additionally, Benghal dayflower has short hairs (pubescence) on the upper leaf surface, unlike Virginia dayflower which is typically scabrous.

Victor Maddox, PhD

MSU Geosystems Research Institute

Box 9555, Mississippi State, MS 39762-9555

Phone (662)325-2313, E-mail: vmaddox@gri.msstate.edu

www.gri.msstate.edu

