

**Plant Community Response in Small Plots One Year after Treatment with
Triclopyr and Endothall in Noxon Rapids Reservoir, MT, 2011**



Ryan M. Wersal and John D. Madsen

Geosystems Research Institute, Mississippi State University, Mississippi State, MS

A Report Submitted to Sanders County, MT, the Montana Weed Trust,
and the U.S. Army Engineer Research and Development Center

November 2011

Geosystems Research Institute Report #5049



MISSISSIPPI STATE
UNIVERSITY™



Preface

This report presents data collected by the Geosystems Research Institute, Mississippi State University in 2010 and 2011 on Noxon Rapids Reservoir. Funding was provided by a grant under the American Recovery and Re-Investment Act and the Montana Weed Trust as a subcontract from Sanders County, Montana to Mississippi State University. We also thank Celestine Duncan, Brian Burke, Heidi Sedivy, and John Halpop for assistance with planning and on the ground logistics. Field assistance was provided by Dr. Wade Givens, Dr. Eric Dibble, Amanda Fernandez, Jonathan Fleming, and Cheryl McLaurin, all from Mississippi State University. Any errors in presentation or fact, however, are the responsibility of the authors.

Plant Community Response in Small Plots One Year after Treatment with Triclopyr and Endothall in Noxon Rapids Reservoir, MT, 2011

Ryan M. Wersal and John D. Madsen

Geosystems Research Institute, Mississippi State University, Mississippi State, MS

Introduction

Aquatic plants are important to lake ecosystems (Madsen et al. 1996, Wetzel 2001) and are essential in promoting the diversity and function of an aquatic system (Carpenter and Lodge 1986). Littoral zone habitat and associated plants may be responsible for a significant proportion of primary production for the entire lake (Ozimek et al. 1990, Wetzel 2001). Littoral zone habitats are prime areas for the spawning of most fish species, including many species important to sport fisheries (Savino and Stein 1989). Furthermore, aquatic plants anchor soft sediments, stabilize underwater slopes, remove suspended particles, and remove nutrients from overlying waters (Barko et al. 1986, Doyle 2000, Madsen et al. 2001). The introduction of non-native plants into littoral zone habitats often alters the complex interactions occurring in these areas (Madsen 1998). Dense stands of non-native plants are often responsible for reduction in oxygen exchange, depletion of dissolved oxygen, increases in water temperatures, and internal nutrient loading (Madsen 1998).

Eurasian watermilfoil (*Myriophyllum spicatum* L.) is a non-native invasive species that, when present, has been associated with declines in native plant species richness and diversity (Madsen et al. 1991, Madsen et al. 2008). Eurasian watermilfoil also poses nuisance problems to humans in the form of increasing flood frequency and intensity, impeding navigation, and limiting recreation opportunities (Madsen et al. 1991). Furthermore, the establishment of Eurasian watermilfoil and subsequent spread is likely perpetuated by the ease of fragmentation (both physical and physiological) of this plant, water movement within the reservoir, and high watercraft traffic that moves fragments to new areas.

Although the impacts from Eurasian watermilfoil are numerous, controlling this species is often difficult and unpredictable. Flowing water, such as the Lower Clark Fork River, further complicates the use of herbicides as water flow will increase the dilution and dissipation of the herbicides. Herbicide applications in run of the river reservoirs are often subject to more extreme perturbations than those of natural lakes. Run of the river reservoirs have variable water-exchange patterns, typically tied to dam operations, which will impact aqueous distribution of herbicides resulting in reduced chemical exposure times against target plants and unacceptable effectiveness (Getsinger et al. 1997).

The use of auxin mimicking herbicides such as 2,4-D and triclopyr, and the contact herbicide endothall have been used extensively for Eurasian watermilfoil control. Additionally, herbicide concentration exposure time (CET) relationships have been designed under controlled conditions to guide management decisions on choosing the correct herbicide concentration with respect to contact time (Netherland et al. 1991, Netherland and Getsinger 1992). However, little data exists with respect to combining a contact herbicide with a systemic herbicide to reduce the exposure

time requirements and maintain plant control. Mesocosm trials of herbicide combinations resulted in reduced contact time needed for effective Eurasian watermilfoil control, while maintaining the benefits of the longer term control afforded by the systemic herbicide (Madsen et al. 2010). Though small scale trials have been conducted, there has been limited field assessment of this herbicide combination. Pursuant to this, effective herbicide concentrations used in field situations still need to be determined; and the selectivity spectrum of non-target plants to this combination is still unknown. Therefore, our objectives of this study were to:

- 1) Demonstrate at the field scale the effectiveness of combining triclopyr with endothall for control of Eurasian watermilfoil and curlyleaf pondweed in flowing systems.
- 2) Evaluate the aquatic plant community response to herbicide treatments one year after treatment.

Materials and Methods

Point Intercept Assessments. Pretreatment (0 weeks after treatment (WAT)) point intercept surveys were conducted on July 20, 2010 using a 50 m grid to assess the plant community in four plots on Noxon Rapids Reservoir prior to herbicide application. The four plots selected were based on surveys conducted throughout the reservoir in 2008 and 2009 (Madsen and Cheshier 2009, Wersal et al. 2009, Wersal et al. 2010a). Plots 2 (23.8 acres) and 4 (28.5 acres) served as our untreated reference plots, meaning no herbicides were placed in these plots. Plot 7 (28.3 acres) was treated with triclopyr alone, and plot 8 (15.8 acres) was treated with the combination of triclopyr + endothall. Additional surveys of each plot occurred at 7 WAT and 52 WAT to assess herbicide efficacy on Eurasian watermilfoil and curlyleaf pondweed; as well as to assess non-target effects on the entire aquatic plant community.

Survey methods were similar to those utilized during recent projects in the Pacific Northwest (Madsen and Wersal 2008, Madsen and Wersal 2009, Wersal et al. 2010a, Wersal et al. 2010b). A total of 36, 38, 35, and 37 points were surveyed in Plots 2, 4, 7, and 8 respectively. Surveys were conducted by boat using Global Positioning System (GPS) technology to navigate to each point. Survey accuracy was 1-3 m (3-10 ft) depending on satellite reception. At each survey point, a weighted thatch rake was deployed to determine the presence of plant species. Spatial data were recorded electronically using FarmWorks Site Mate[®] software (Hamilton, IN). The software allowed for in-field geographic and attribute data collection. Data were recorded in database templates using specific pick lists constructed exclusively for this project. Site Mate[®] provided an environment for displaying geographic and attribute data and enabled navigation to specific locations on the lake.

Statistical Analyses. Plant species presence was averaged over all points sampled and multiplied by 100 to calculate percent frequency. Changes in the occurrence of plant species between the pretreatment survey and 7 WAT and 52 WAT surveys were determined using the McNemar's test. The McNemar's test is used to assess the differences in the correlated proportions within a given data set between variables that are not independent, i.e. sampling the same points pre- and post-treatment (Stokes et al. 2000, Wersal et al. 2006, Wersal et al. 2010b).

All comparisons were made back to pretreatment plant occurrence. Mean species richness, native species richness, and non-native species richness was calculated for each plot and subjected to a general linear model. If a significant difference in species richness was detected, means were separated using a Fisher's Protected LSD test. All analyses were conducted at a $p < 0.05$ significance level.

Results and Discussion

Plot 7 (Triclopyr Only)

The presence of Eurasian watermilfoil in Plot 7 significantly declined from 50% before herbicide treatment to 16% 7 WAT and 12% 52 WAT (Table 1 and Figure 1). This represents 70% and 76% control respectively for the 7 and 52 WAT surveys. Curlyleaf pondweed was observed at 81%, 50%, and 67% of survey points during the pre, 7, and 52 WAT surveys respectively (Figure 2). The decline in plot 7 is most likely due to natural senescence (Woolf and Madsen 2003), as plants had produced turions and were falling out of the water column during the time of the 7 WAT survey. In the spring of 2011, the lower Clark Fork River systems experienced a higher than average snow-melt resulting in high water conditions for most of the summer. The high water reduced light penetration and likely kept water temperature below normal. These factors likely resulted in delayed germination and sprouting of most plant species and thus during the 52 WAT survey curlyleaf pondweed was still actively growing in more locations in the plot.

No significant impact was observed from the herbicide application on native plant species at either the 7 WAT or 52 WAT surveys. Species richness and native species richness were not different between survey times as well. The presence of elodea in this plot increased from the pretreatment survey to the 52 WAT survey, suggesting that as Eurasian watermilfoil was removed Elodea was able to re-colonize those areas. Similar results were observed in Hayden Lake, ID when applications of 2,4-D and triclopyr were made for Eurasian watermilfoil control (Wersal et al. 2010b). The use of triclopyr alone resulted in very selective control of Eurasian watermilfoil for at least a full year after treatment, a result due to the CET achieved after treatment (Wersal and Madsen 2011). A longer exposure time would have likely resulted in non-target plant injury, especially to northern watermilfoil (*Myriophyllum sibiricum*) and white water-buttercup (*Ranunculus aquatilis*).

Plot 8 (Triclopyr and Endothall)

Eurasian watermilfoil in plot 8 significantly declined by 7 WAT and control was maintained to 52 WAT with the combination of triclopyr + endothall (Table 2). Eurasian watermilfoil was observed at 63%, 9%, and 5% of survey points during the pretreatment, 7 WAT, and 52 WAT surveys respectively (Figure 3). These results represent 92% reduction in Eurasian watermilfoil occurrence out to 52 WAT. The presence of curlyleaf pondweed also significantly declined from 74% pretreatment to 3% 7 WAT (Figure 4). Curlyleaf pondweed was found at 43% of the survey points at 52 WAT. The decline at 7 WAT was likely due to the endothall applied in the plot. Endothall, being a contact herbicide, would have had an immediate effect on plants in the year of treatment. Curlyleaf pondweed has begun to recover as indicated by the 52 WAT survey.

The combination of triclopyr + endothall was much less selective in plot 8, as there were impacts to the native plant community. All species richness metrics for both the 7 and 52 WAT surveys were lower than what was estimated during the pretreatment surveys. The presence of elodea increased during the 52 WAT survey, again likely due to opening new areas for expansion and growth by the herbicide application. Although, the combination herbicide treatment was effective there will be a trade-off in selectivity when compared to using triclopyr alone. Given the estimated half-life for dye and herbicide in plot 8 (Wersal and Madsen 2011), triclopyr applied alone would have been much less effective. In areas where there is potential for high water exchange, the combination treatment would be necessary to maximize control.

Plots 2 and 4 (Untreated Reference Plots)

The plant community in plot 2 has changed little over the course of the study. Eurasian watermilfoil was found at survey points during the post treatment surveys where it was not observed during previous surveys, suggesting that the population in plot 2 is expanding, though statistical differences in presence have not been detected (Table 3). Species richness did not change between any of the surveys. The locations of Eurasian watermilfoil and curlyleaf pondweed for plot 2 are depicted in figures 5 and 6 respectively.

The presence of Eurasian watermilfoil in plot 4 did not change between the pretreatment survey and the 7 and WAT surveys (Table 4). Therefore, the reductions observed in Eurasian watermilfoil in plots 7 and 8 can be attributed to the herbicide applications. The locations of Eurasian watermilfoil and curlyleaf pondweed for plot 4 are depicted in figures 7 and 8 respectively.

Conclusions. Herbicide applications were effective at reducing the presence of Eurasian watermilfoil in the treated plots, 76% and 92% for plots 7 and 8 respectively to 52 WAT. Control of Eurasian watermilfoil is achievable in flowing water systems if there is an understanding of water exchange characteristics at a given site. Water exchange is likely to be site specific within Noxon so additional studies are needed, especially upstream, to develop a water exchange data set for portions of the reservoir to base management decisions on.

Our data indicate that Eurasian watermilfoil can be selectively removed from areas of Noxon, and that native species will rapidly re-colonize areas once inhabited by Eurasian watermilfoil. Furthermore, these data suggest that Eurasian watermilfoil control can be maintained for at least two growing seasons with a single herbicide application. Achieving multiple year control would allow for the treatment of additional areas without having to continually re-treat in the same plots. Though this will depend upon site location, water flow, and distance from other Eurasian watermilfoil infestations that would re-colonize an already treated area.

The combination herbicide treatment was less selective than applying triclopyr alone. Using triclopyr alone would not have been effective in plot 8, as the necessary exposure time would not have been met to achieve acceptable results. Therefore, the use of triclopyr alone will not be conducive to all places in the reservoir, especially in areas of increased water-exchange; these areas will need the combination treatment to meet CET requirements. The potential short term impacts of herbicide applications on the native plant community should not overshadow the

long-term effects that Eurasian watermilfoil will have if left unmanaged. Species such as leafy pondweed (*Potamogeton foliosus*) and elodea which are widespread in Noxon recovered by 52 WAT to levels similar or greater than what was observed during the pretreatment survey in plot 8. There is a native propagule bank present in Noxon that will allow the native community to recover following Eurasian watermilfoil management.

Literature Cited

- Barko, J.W., M.S. Adams, and N.S. Clesceri. 1986. Environmental factors and their consideration in the management of submersed aquatic vegetation: A review. *Journal of Aquatic Plant Management* 24:1-10.
- Doyle, R.D., 2000. Effects of navigation on aquatic plants: effects of sediment resuspension and deposition on plant growth and reproduction. Upper Mississippi River – Illinois Waterway System Navigation Study, ENV Rep. 28. U.S. Army Corps of Engineers, Rock Island District, St. Louis District, St. Paul District. 64 pp.
- Getsinger, K.D., E.G. Turner, J.D. Madsen, and M.D. Netherland. 1997. Restoring native vegetation in a Eurasian watermilfoil dominated plant community using the herbicide triclopyr. *Regulated Rivers: Research and Management* 13:357-375.
- Madsen, J.D. 1998. Predicting invasion success of Eurasian watermilfoil. *Journal of Aquatic Plant Management* 36:28-32.
- Madsen, J.D., J.W. Sutherland, J.A. Bloomfield, L.W. Eichler, and C.W. Boylen. 1991. The decline of native vegetation under dense Eurasian watermilfoil canopies. *Journal of Aquatic Plant Management* 29:94-99.
- Madsen, J.D., J.A. Bloomfield, J.W. Sutherland, L.W. Eichler, and C.W. Boylen. 1996. The aquatic macrophyte community of Onondaga Lake: Field survey and plant growth bioassays of lake sediments. *Lake and Reservoir Management* 12:73-79.
- Madsen, J.D., P.A. Chambers, W.F. James, E.W. Koch, and D.F. Westlake. 2001. The interactions between water movement, sediment dynamics and submersed macrophytes. *Hydrobiologia* 444:71-84.
- Madsen, J.D. and R.M. Wersal. 2008. Assessment of Eurasian watermilfoil (*Myriophyllum spicatum* L.) populations in Lake Pend Oreille, Idaho for 2007. GeoResources Institute Report 5028, 116p.
- Madsen, J.D., R.M. Stewart, K.D. Getsinger, R.L. Johnson, and R.M. Wersal. 2008. Aquatic plant communities in Waneta Lake and Lamoka Lake, New York. *Northeastern Naturalist* 15:97-110.

- Madsen, J. D. and J. C. Cheshier. 2009. Eurasian watermilfoil survey of three reservoirs in the Lower Clark Fork River, Montana: I. Results of the field vegetation survey. Geosystems Research Institute Report 5033, 59p.
- Madsen, J.D. and R.M. Wersal. 2009. Aquatic plant community and Eurasian watermilfoil (*Myriophyllum spicatum* L.) management assessment in Lake Pend Oreille, Idaho for 2008. Geosystems Research Institute Report 5032, 65p.
- Madsen, J.D., R.M. Wersal, K.D. Getsinger, and J.G. Skogerboe. 2010. Combinations of endothall with 2,4-D and triclopyr for Eurasian watermilfoil control. APCRP Technical Notes Collection (ERDC/TN APCRP-CC-14). U.S. Army Engineer Research and Development Center, Vicksburg, MS. 10p.
- Netherland, M.D., W.R. Green, and K.D. Getsinger. 1991. Endothall concentration and exposure time relationships for the control of Eurasian watermilfoil and hydrilla. *Journal of Aquatic Plant Management* 29:61-67.
- Netherland, M.D. and K.D. Getsinger. 1992. Efficacy of triclopyr on Eurasian watermilfoil: concentration and exposure time effects. *Journal of Aquatic Plant Management* 30:1-5.
- Ozimek, T., R.D. Gulati, and E. van Donk. 1990. Can macrophytes be useful in biomanipulation of lakes? The Lake Zwemlust example. *Hydrobiologia* 200/201:399-407.
- Savino, J.F. and R.A. Stein. 1989. Behavior of fish predators and their prey: habitat choice between open water and dense vegetation. *Environmental Biology of Fishes* 24:287-293.
- Stokes, M.E., C.S. Davis, and G.G. Koch. 2000. *Categorical Data Analysis Using the SAS[®] System*, second edition. SAS Institute Inc., Cary, NC, USA.
- Wersal, R.M., J.D. Madsen, B.R. McMillan and P.D. Gerard. 2006. Environmental factors affecting biomass and distribution of *Stuckenia pectinata* in the Heron Lake System, Minnesota, USA. *Wetlands* 26:313-321.
- Wersal, R.M., J.D. Madsen, and J.C. Cheshier. 2009. Eurasian watermilfoil monitoring and mapping in Noxon Rapids Reservoir in 2009. Geosystems Research Institute Report 5041, 11p.
- Wersal, R.M., J.D. Madsen, and J.C. Cheshier. 2010a. Aquatic plant monitoring in Noxon Rapids Reservoir and Cabinet Gorge Reservoir for 2010. Geosystems Research Institute Report 5042, 18p.
- Wersal, R.M., J.D. Madsen, T.E. Woolf, and N. Eckberg. 2010b. Assessment of herbicide efficacy on Eurasian watermilfoil and impacts to the native submersed plant community in Hayden Lake, Idaho, USA. *Journal of Aquatic Plant Management* 48:5-11.

- Wersal, R.M. and J.D. Madsen. 2011. Evaluating plant response to triclopyr applied alone and in combination with endothall in Noxon Rapids Reservoir for 2010: Phase 2. Geosystems Research Institute Report 5046, 64p.
- Wetzel, R.G. 2001. Limnology: Lake and River Ecosystems, Third Edition. Academic Press, San Diego, CA, USA. 1006 pp.
- Woolf, T.E. and J.D. Madsen. 2003. Seasonal biomass and carbohydrate allocation patterns in southern Minnesota curlyleaf pondweed populations. *Journal of Aquatic Plant Management* 41:113-118.

Table 1. Aquatic plant occurrence in plot 7 herbicide (triclopyr alone) treatment area in Noxon Rapids Reservoir, MT 2010-2011. Differences between sampling events were determined at a $p < 0.05$ significance level using the McNemars test. An asterisk indicates a significant change from the pretreatment occurrence for each species. Mean species richness ($\pm 1SE$) data were separated using the LSD method, values within a row sharing the same letter are not different at $p < 0.05$ significance level.

Plant Species	Common Name	0 WAT	7 WAT	52 WAT
		% Occurrence	% Occurrence	% Occurrence
<i>Ceratophyllum demersum</i>	Coontail	38	44	61
<i>Chara sp.</i>	Muskgrass	16	19	21
<i>Elodea canadensis</i>	Elodea	47	47	73*
<i>Heteranthera dubia</i>	Water stargrass	9	6	6
<i>Myriophyllum sibiricum</i>	Northern watermilfoil	38	22	24
<i>Myriophyllum spicatum</i>	Eurasian watermilfoil	50	16*	12*
<i>Potamogeton crispus</i>	Curlyleaf pondweed	81	50*	67
<i>Potamogeton foliosus</i>	Leafy pondweed	31	16	42
<i>Potamogeton illinoensis</i>	Illinois pondweed	0	3	0
<i>Potamogeton richardsonii</i>	Clasping-leaved pondweed	13	22	24
<i>Ranunculus aquatilis</i>	White water-buttercup	22	19	24
<i>Stuckenia pectinata</i>	Sago pondweed	59	50	42
Species Richness		3.8 \pm 0.3a	3.0 \pm 0.3b	3.9 \pm 0.3a
Native Richness		2.6 \pm 0.2a	2.4 \pm 0.2a	3.2 \pm 0.3a
Non-native Richness		1.2 \pm 0.1a	0.6 \pm 0.1b	0.8 \pm 0.1b

Table 2. Aquatic plant occurrence in plot 8 herbicide (triclopyr + endothall) treatment area in Noxon Rapids Reservoir, MT 2010-2011. Differences between sampling events were determined at a $p < 0.05$ significance level using the McNemars test. An asterisk indicates a significant change from the pretreatment occurrence for each species. Mean species richness ($\pm 1SE$) data were separated using the LSD method, values within a row sharing the same letter are not different at $p < 0.05$ significance level.

Plant Species	Common Name	0 WAT	7 WAT	52 WAT
		% Occurrence	% Occurrence	% Occurrence
<i>Ceratophyllum demersum</i>	Coontail	57	37	30
<i>Chara sp.</i>	Muskgrass	9	37*	8
<i>Elodea canadensis</i>	Elodea	31	51	65*
<i>Heteranthera dubia</i>	Water stargrass	3	3	5
<i>Myriophyllum sibiricum</i>	Northern watermilfoil	40	3*	8*
<i>Myriophyllum spicatum</i>	Eurasian watermilfoil	63	9*	5*
<i>Potamogeton crispus</i>	Curlyleaf pondweed	74	3*	43*
<i>Potamogeton foliosus</i>	Leafy pondweed	11	0*	19
<i>Potamogeton illinoensis</i>	Illinois pondweed	3	0	0
<i>Potamogeton richardsonii</i>	Clasping-leaved pondweed	31	3*	3*
<i>Ranunculus aquatilis</i>	White water-buttercup	46	20	22
<i>Stuckenia pectinata</i>	Sago pondweed	40	3*	19
<i>Vallisneria americana</i>	Wildcelery	0	6	0
Species Richness		3.9 \pm 0.3a	1.7 \pm 0.2b	2.4 \pm 0.3b
Native Richness		2.5 \pm 0.3a	1.6 \pm 0.2b	2.0 \pm 0.3b
Non-native Richness		1.3 \pm 0.1a	0.1 \pm 0.1b	0.5 \pm 0.1c

Table 3. Aquatic plant occurrence in plot 2 untreated reference area in Noxon Rapids Reservoir, MT 2010-2011. Differences between sampling events were determined at a $p < 0.05$ significance level using the McNemars test. An asterisk indicates a significant change from the pretreatment occurrence for each species. Mean species richness ($\pm 1SE$) data were separated using the LSD method, values within a row sharing the same letter are not different at $p < 0.05$ significance level.

Plant Species	Common Name	0 WAT	7 WAT	52 WAT
		% Occurrence	% Occurrence	% Occurrence
<i>Butomus umbellatus</i>	Flowering rush	0	0	3
<i>Ceratophyllum demersum</i>	Coontail	61	53	61
<i>Chara sp.</i>	Muskgrass	6	14	19
<i>Elodea canadensis</i>	Elodea	50	75*	70
<i>Heteranthera dubia</i>	Water stargrass	11	8	17
<i>Myriophyllum sibiricum</i>	Northern watermilfoil	11	0*	6
<i>Myriophyllum spicatum</i>	Eurasian watermilfoil	67	80	86
<i>Najas flexilis</i>	Slender naiad	0	0	3
<i>Potamogeton crispus</i>	Curlyleaf pondweed	75	61	55*
<i>Potamogeton foliosus</i>	Leafy pondweed	31	22	58*
<i>Potamogeton illinoensis</i>	Illinois pondweed	17	14	3
<i>Potamogeton praelongus</i>	Whitestem pondweed	3	3	5
<i>Potamogeton richardsonii</i>	Clasping-leaved pondweed	31	17	24
<i>Ranunculus aquatilis</i>	White water-buttercup	6	6	3
<i>Stuckenia pectinata</i>	Sago pondweed	31	58*	29
Species Richness		3.9 \pm 0.3a	4.1 \pm 0.3a	4.5 \pm 0.3a
Native Richness		2.6 \pm 0.2a	2.7 \pm 0.2a	3.1 \pm 0.3a
Non-native Richness		1.4 \pm 0.1a	1.4 \pm 0.1a	1.5 \pm 0.1a

Table 4. Aquatic plant occurrence in plot 4 untreated reference area in Noxon Rapids Reservoir, MT 2010-2011. Differences between sampling events were determined at a $p < 0.05$ significance level using the McNemars test. An asterisk indicates a significant change from the pretreatment occurrence for each species. Mean species richness ($\pm 1SE$) data were separated using the LSD method, values within a row sharing the same letter are not different at $p < 0.05$ significance level.

Plant Species	Common Name	0 WAT	7 WAT	52 WAT
		% Occurrence	% Occurrence	% Occurrence
<i>Butomus umbellatus</i>	Flowering rush	7	0*	0*
<i>Ceratophyllum demersum</i>	Coontail	50	67	50
<i>Chara sp.</i>	Muskgrass	7	0*	3
<i>Elodea canadensis</i>	Elodea	30	40	53
<i>Heteranthera dubia</i>	Water stargrass	3	10	17
<i>Myriophyllum sibiricum</i>	Northern watermilfoil	33	40	33
<i>Myriophyllum spicatum</i>	Eurasian watermilfoil	30	27	20
<i>Nitella sp.</i>	Nitella	0	13*	0
<i>Potamogeton crispus</i>	Curlyleaf pondweed	46	3*	20*
<i>Potamogeton foliosus</i>	Leafy pondweed	13	30	37
<i>Potamogeton illinoensis</i>	Illinois pondweed	3	0	3
<i>Potamogeton richardsonii</i>	Clasping-leaved pondweed	20	10	20
<i>Ranunculus aquatilis</i>	White water-buttercup	20	10	13
<i>Stuckenia pectinata</i>	Sago pondweed	23	17	27
Species Richness		2.7 \pm 0.4a	2.7 \pm 0.3a	2.9 \pm 0.4a
Native Richness		2.0 \pm 0.3a	2.3 \pm 0.3a	2.6 \pm 0.4a
Non-native Richness		0.8 \pm 0.1a	0.3 \pm 0.1b	0.4 \pm 0.1b




Figure 1. Locations of Eurasian watermilfoil pretreatment, 7 weeks after treatment, and 52 weeks after treatment in plot 7, which received an application of triclopyr alone.




Figure 2. Locations of curlyleaf pondweed pretreatment, 7 weeks after treatment, and 52 weeks after treatment in plot 7, which received an application of triclopyr alone.




Figure 3. Locations of Eurasian watermilfoil pretreatment, 7 weeks after treatment, and 52 weeks after treatment in plot 8, which received an application of triclopyr + endothall.




Figure 4. Locations of curlyleaf pondweed pretreatment, 7 weeks after treatment, and 52 weeks after treatment in plot 8, which received an application of triclopyr + endothall.




Figure 5. Locations of Eurasian watermilfoil pretreatment, 7 weeks after treatment, and 52 weeks after treatment in plot 2, which served as an untreated reference plot.




Figure 6. Locations of curlyleaf pondweed pretreatment, 7 weeks after treatment, and 52 weeks after treatment in plot 2, which served as an untreated reference plot.




Figure 7. Locations of Eurasian watermilfoil pretreatment, 7 weeks after treatment, and 52 weeks after treatment in plot 4, which served as an untreated reference plot.




Figure 8. Locations of curlyleaf pondweed pretreatment, 7 weeks after treatment, and 52 weeks after treatment in plot 4, which served as an untreated reference plot.