

Invasion Alert

Cite this article: Gebhart MG, Schmid SA, Turner S, Webb D, Thum R, Beets J, Turnage S (2025) Invasive eelgrass hybrid (*Vallisneria* × *pseudorosulata*) in the southeastern United States. *Invasive Plant Sci. Manag.* **18**(e2), 1–5. doi: [10.1017/inp.2024.37](https://doi.org/10.1017/inp.2024.37)

Received: 23 September 2024

Revised: 8 November 2024

Accepted: 15 November 2024

Associate Editor:

Ryan M. Wersal, Minnesota State University

Keywords:

aquarium trade; aquatic macrophyte; cryptic invader; submersed; hybrid

Corresponding author:

Maxwell G. Gebhart;

Email: mgebhart@gri.msstate.edu

Invasive eelgrass hybrid (*Vallisneria* × *pseudorosulata*) in the southeastern United States

Maxwell G. Gebhart¹ , Samuel A. Schmid¹ , Stephen Turner², David Webb³, Ryan Thum⁴, Jens Beets⁵ and Gray Turnage⁶

¹Research Associate, Mississippi State University - Geosystems Research Institute, Starkville, MS, USA; ²Program Manager, Tennessee Valley Authority, Knoxville, TN, USA; ³Senior Program Manager, Tennessee Valley Authority, Knoxville, TN, USA; ⁴Professor, Montana State University, Bozeman, MT, USA; ⁵Research Ecologist, USDA, Davis, CA, USA and ⁶Extension Research Professor, Mississippi State University - Geosystems Research Institute, Starkville, MS, USA

Abstract

Vallisneria × *pseudorosulata* S. Fujii & M. Maki is an invasive aquatic weed that has recently become a major issue within the U.S. Southeast. *Vallisneria* × *pseudorosulata* is a hybrid between two nonnative eelgrass species (*Vallisneria spiralis* L. and *Vallisneria denseserrulata* Makino) and has rapidly overtaken water bodies in Tennessee, Alabama, and Florida. This hybrid can reproduce rapidly through offshoot formation and floating propagules capable of drifting large distances before establishing. *Vallisneria* × *pseudorosulata* has been previously found in Japan and is thought to have been introduced in the United States by the aquarium trade or through dumping.

Vallisneria L. is a genus of submersed macrophytes that have often been a major target for phytoremediation and restoration efforts (Cao and Ruan 2015; Korschgen and Green 1988; Les et al. 2008). Often, American eelgrass or *Vallisneria americana* Michx. is a primary species for restoration efforts in the United States due to the natural habitat, forage, and structure it can provide to many ecosystems; however, there has been a steady decline of the species throughout many aquatic systems (Engelhardt et al. 2014). Simultaneously, recent research has determined that there are two native species of *Vallisneria* in the United States, *V. americana* and *Vallisneria neotropicalis* Vict. (southern eelgrass; Les et al. 2008; Martin and Mort 2023). Thus, most of the current research on *Vallisneria* has focused on how to restore *V. americana* to the landscape and reasons why it may be disappearing. However, there has been a sudden and rapid invasion by shortstemmed eelgrass, *V. × pseudorosulata* S. Fujii & M. Maki (= *Vallisneria spiralis* L. [tapegrass] × *Vallisneria denseserrulata* Makino [toothed eelgrass]) which has received increasing attention in the past 5 yr. *Vallisneria* × *pseudorosulata* is thought to have been introduced from the aquarium trade as an ornamental *Vallisneria* plant for sale but was later dumped into a water body and has spread rapidly throughout the U.S. Southeast (Gorham et al. 2021; Wasekura et al. 2016). It is also thought to have been used in native *Vallisneria* restoration under the guise of being a native hybrid (King's Bay Restoration Project, Crystal River, FL). Both parental taxa of *V. × pseudorosulata*, *V. spiralis* and *V. denseserrulata*, are nonnative to the United States and originate from the Eurasian continent (Chen et al. 2012; Gorham et al. 2021; Les et al. 2008; Mesterházy et al. 2021; Wasekura et al. 2016). *Vallisneria* × *pseudorosulata* has infested the Tennessee Valley Authority (TVA) system; this population is thought to have established sometime between 2018 and 2019 (Gorham et al. 2021).

The introduction into the TVA system is not an isolated event, as *V. × pseudorosulata* has been found in multiple systems in Florida, Alabama, Tennessee, and recently Mississippi. Internationally, *V. × pseudorosulata* was initially reported in 2016 as present in multiple water bodies throughout Japan (Wasekura et al. 2016). The cross is unlikely to have occurred naturally, as both parent species are completely disjunct (Gorham et al. 2021; Wasekura et al. 2016). Since this hybridization, *V. × pseudorosulata* has been distributed widely through the aquarium trade under the names *V. spiralis*, *Vallisneria* 'Rock Star', and potentially other common names (Martin and Mort 2023; Padilla and Williams 2004; Wasekura et al. 2016). Gorham et al. (2021) confirmed *V. × pseudorosulata* primarily in Florida, with one site in Alabama, the present report confirms several other sites where *V. × pseudorosulata* is found in the U.S. Southeast. There is little understanding about *V. × pseudorosulata* and its ecological interactions with other organisms. However, there is anecdotal evidence that *V. × pseudorosulata* is rapidly displacing the invasive species hydrilla [*Hydrilla verticillata* (L. f.) Royle], prompting major concern from resource managers (Gorham et al. 2021; Wetzel 2020). Whether or not *H. verticillata* is being displaced by *V. × pseudorosulata*, it is incredibly clear that *V. × pseudorosulata* can rapidly

© The Author(s), 2024. Published by Cambridge University Press on behalf of Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



Table 1. Collection and location details of *Vallisneria × pseudorosulata* collected in the United States in 2023 and 2024^a.

Primary collector	Collection no.	Catalog no.	Date	State	Latitude °N	Longitude °W
ST	1	MISSA039774	September 2023	Alabama	34.6421	85.9696
ST	2	MISSA039775	September 2023	Tennessee	35.0634	85.5323
ST	3	MISSA039769	September 2023	Alabama	34.5601	86.8517
ST	4	MISSA039770	September 2023	Tennessee	35.1561	85.1559
SAS	353	MISSA039771	June 24, 2024	Mississippi	34.4758	88.3346

^aAll specimens have been submitted and digitized in the Mississippi State University herbarium (MISSA).




Figure 1. (A) The typical growth form of *Vallisneria × pseudorosulata* as a rosette with long, thin leaves. (B) Asexual reproduction usually takes form of stolons that extend from the rosette base. (C) When established, stolons with many small ramets can extend more than 30 cm. (D) *V. × pseudorosulata* leaves are denoted by a bright green lacunal band that follows the midvein. (E) Leaf tips for *V. × pseudorosulata* are obtusely angled and appear rounded. (F) Serrations are small and dense along leaf margins. (G) Stem formation by *V. × pseudorosulata* during experimentation in mesocosms. (H) Chains of ramets can form in both still and moving water, which can allow for new establishment once the ramets sink. (I) Pistillate flowers of *V. × pseudorosulata*, which are the only flower type currently found.

become the dominant plant within a water body (Gorham et al. 2021). It is currently hypothesized that *V. × pseudorosulata* grows as an evergreen perennial and can form large floating mats of propagules, upward of 16 km (10 miles), which has led to its rapid spread (Anonymous 2023; Plotka 2023; Sapp 2024). The

propagules are capable of floating for long stretches and may be why Mississippi was recently invaded by *V. × pseudorosulata*. The population of *V. × pseudorosulata* found in Mississippi was discovered in June 2024 in Pool E (Prentiss County, near Belmont) of the Tennessee-Tombigbee Waterway, a system that is connected




Figure 2. (A) A comparison between *Vallisneria* × *pseudorosulata* (left), *Vallisneria americana* (center), and *Vallisneria neotropalis* (right). (B) A comparison of the leaf colorations between *V.* × *pseudorosulata* (left), *V. americana* (center), and *V. neotropalis* (right). Both *V. americana* and *V. neotropalis* leaves can have different colorations based on the growing depth, but may be typically darker than *V.* × *pseudorosulata* leaves.




Figure 3. A map displaying known locations of *Vallisneria* × *pseudorosulata* in the U.S. Southeast. The circles represent approximate sampling locations from Gorham et al. (2021), and the squares represent the population extent being reported here.

to the TVA system, flows into Mobile River, and ultimately drains into the Gulf of Mexico at Mobile Bay (Schmid and Magandy 353 MISSA039771; Table 1). This is the first report to document the newest population of *V.* × *pseudorosulata* as well as the distance of spread from the sampling location Gorham et al. (2021) reported in Lake Guntersville.

Morphologically, many *Vallisneria* species are incredibly similar with few diagnostic characteristics; this crypticity equally applies to *V.* × *pseudorosulata*, which can impede early detection (Les et al. 2008). *Vallisneria* × *pseudorosulata* looks similar to *V. americana*, growing as a rosette with long, ribbon-like leaves and reduced floral characteristics that can make in-field identification

difficult (Les et al. 2008; Martin and Mort 2023). Often, *V. × pseudorosulata* can be identified by a bright green lacunal band running up the midvein to the leaf tip; however, this can be difficult to determine on young leaves or if water clarity is an issue (Figures 1D and 2). Another characteristic, and the most diagnostic, is a partially lignified stem that *V. × pseudorosulata* will grow just above the soil line (Wasekura et al. 2016; MGG, personal observation). This stem structure was described by Wasekura et al. (2016) that grows on mature plants, but not every *V. × pseudorosulata* individual has shown this in the field (MGG, personal observation). This stem can be 1 to 12 cm in length and typically has small, but apparent internodes with adventitious roots growing from the nodes (Figure 1G and 1H). Currently it is highly recommended that if a *Vallisneria* population should look similar to *V. × pseudorosulata*, then genetic testing should be done for confirmation. Genetic testing is highly accurate, and assays have been developed to distinguish *V. × pseudorosulata* from *V. americana* and *V. neotropicalis* (Martin and Mort 2023; Tringali et al. 2023). Many *Vallisneria* species typically reproduce via runners that have been observed growing above and below the soil line, often with upward of 10 to 20 daughter plants (Korschgen and Green 1988; Martin and Mort 2023; McFarland and Shafer 2008). Anecdotally, these runners have been observed detaching from the parent plant and floating long distances before settling in the sediment again. *Vallisneria* spp. are dioecious, but *V. × pseudorosulata* has only been observed with pistillate flowers, leading researchers to hypothesize it does not reproduce sexually (Gorham et al. 2021; Martin and Mort 2023; Figure 1I). Preliminary observations have also shown that *V. × pseudorosulata* can handle low to freezing temperatures, allowing rapid establishment in the spring (MGG, personal observation). *Vallisneria × pseudorosulata* presents not only a unique challenge in the field with identification, but the entire genus of *Vallisneria* has presented multiple complexities for systematists, taxonomists, and evolutionary biologists. Particularly, species like ribbonweed, *Vallisneria australis* S.W.L. Jacobs & Les and *V. neotropicalis* are still debated in many circles and often get lumped under the name of a congener (Les et al. 2008; Jacobs and Frank 1997).

The evolutionary and taxonomic history of *Vallisneria* is thought to be another major contributor to the confusion surrounding the invasive *V. × pseudorosulata*. Lowden (1982) sought to create a taxonomic tree using floral characteristics specifically, thus leading to only two species of *Vallisneria*, *V. spiralis* and *V. americana*. However, reevaluations of the genus in both 2008 and 2023 determined genetically that there are most likely 12 to 16 species scattered throughout the world (Les et al. 2008; Martin and Mort 2023). In both 2008 and 2023, arguments were made for the distinction of two native *Vallisneria* species in North America, *V. americana* and *V. neotropicalis* (Les et al. 2008; Marie-Victorin 1943; Martin and Mort 2023). *Vallisneria americana* is well understood and has presence throughout the United States. However, there is little knowledge about the range and biology of *V. neotropicalis*. Both native species have been observed co-occurring with *V. × pseudorosulata*, which again can make in situ identification difficult (Figures 2 and 3). These co-occurrences have been genetically confirmed through ITS sequencing with the sampled populations found in Figure 3. We recommend that systems connected to known populations of *V. × pseudorosulata* be monitored for invasion, and when new populations of *Vallisneria* are observed, they should be examined for an elongated stem and, ideally, genetically confirmed. Current invasions have taken over impressively

large areas of the TVA system, most recently in Mississippi, suggesting further spread throughout the Tennessee River system.

The current observations of *V. × pseudorosulata* tied with how little information is available at present raise great concern for further spread throughout the U.S. Southeast. *Vallisneria × pseudorosulata* was first recorded in the United States recently; however, the rate of its invasion and spread is of major concern. Gorham et al. (2021) previously reported only one geographic point in Lake Guntersville where *V. × pseudorosulata* is found; however, our report shows spread both upstream and downstream. Our report is also the first to have visual representations of the structures that make *V. × pseudorosulata* unique for identification and to describe problems with in-field identification. There is a dearth of cohesive research for many of the species in *Vallisneria*, including *V. × pseudorosulata*, therefore, future studies should strive toward creating comprehensive information focused on ecological impacts, biological traits, and management of *V. × pseudorosulata*.

Acknowledgments. We would like to thank Ryan Folk and the rest of the Mississippi State University herbarium (MISSA) team for digitizing and housing voucher specimens. We would like to thank the reviewers for their comments how to improve this invasion alert.

Funding statement. Funding for this work was provided by the Tennessee Valley Authority (TVA agreement no. 13646).

Competing interests. The authors declare no conflicts of interest.

References

- Anonymous (2023) The eelgrass battle: TVA and partners work to wrangle invasive species at Guntersville. QCD News, September 30
- Cao J, Ruan H (2015) Elodeid species as nursery beds for successful seed restoration of *Vallisneria spiralis* L. Pol J Ecol 63(1):53–62
- Chen L, Chen J, Gituru RW, Wang Q (2012) Generic phylogeny, historical biogeography and character evolution of the cosmopolitan aquatic plant family Hydrocharitaceae. BMC Evol Biol 12:30
- Engelhardt KAM, Lloyd MW, Neel MC (2014) Effects of genetic diversity on conservation and restoration potential at individual, population, and regional scales. Biol Conserv 179:6–16
- Gorham SB, Seyoum S, Furman BT, Darnell KM, Reynolds LK, Tringali MD (2021) Molecular detection of a non-native hybrid eelgrass, *Vallisneria spiralis* Linnaeus (1753) × *V. denseserrulata* Makino (1921), in the southeastern United States. Aquat Bot 175:103445
- Jacobs SWL, Frank KA (1997) Notes on *Vallisneria* (Hydrocharitaceae) in Australia, with descriptions of two new species. Telopea 7:111–118
- Korschgen CE, Green WL (1988) American Wildcelery (*Vallisneria americana*): Ecological Considerations for Restoration. Fish and Wildlife Technical Report 19. Washington, DC: U.S. Fish and Wildlife Service. 24 p
- Les DH, Jacobs SWL, Tippery NP, Chen L, Moody ML, Wilstermann-Hildebrand M (2008) Systematics of *Vallisneria* (Hydrocharitaceae). Syst Bot 33:49–65
- Lowden RM (1982) An approach to the taxonomy of *Vallisneria* L. (Hydrocharitaceae). Aquat Bot 13:269–298
- Marie-Victorin F (1943) Les vallisnères américaines. Montreal: Institut botanique de l'Université de Montréal. 21p
- Martin AP, Mort ME (2023) *Vallisneria* (Hydrocharitaceae): novel species, taxonomic revisions, and hybridization. Aquat Bot 188:103669
- McFarland DG, Shafer DJ (2008) Factors influencing reproduction in American wild celery: a synthesis. J Aquat Plant Manag 46:129–144
- Mesterházy A, Somogyi G, Efremov A, Verloove F (2021) Assessing the genuine identity of alien *Vallisneria* (Hydrocharitaceae) species in Europe. Aquat Bot 174:103431

- Padilla DK, Williams SL (2004) Beyond ballast water: aquarium and ornamental trades as sources of invasive species in aquatic ecosystems. *Front Ecol Environ* 2:131–138
- Plotka M (2023) TVA, other agencies tackle eelgrass lake invasion in Guntersville. *WAFF* 48, August 30
- Sapp S (2024) TVA utilizes winter weather to reduce eelgrass species in Guntersville. *WAFF* 48, January 23
- Tringali MD, Gorham SB, Seyoum S, Puchulutegui C, Bass MS, Furman BT, Mallison C (2023) A PCR assay for the detection of introduced *Vallisneria spiralis*, *V. denseserrulata* and their hybrids. *Conserv Genet Resour* 15:125–133
- Wasekura H, Horie S, Fujii S, Maki M (2016) Molecular identification of alien species of *Vallisneria* (Hydrocharitaceae) species in Japan with a special emphasis on the commercially traded accessions and the discovery of hybrid between nonindigenous *V. spiralis* and native *V. denseserrulata*. *Aquat Bot* 128:1–6
- Wetzel M (2020) Homeowners, TVA struggling with aquatic plant life in river. *Gadsden Times*, January 19